


Section 2.4 Dividing Polynomials; Remainder and Factor Theorems

Objectives

- 1 Use long division to divide polynomials.
- 2 Use synthetic division to divide polynomials.
- 3 Evaluate a polynomial using the Remainder Theorem.
- 4 Use the Factor Theorem to solve a polynomial equation.



A moth has moved into your closet. She appeared in your bedroom at night, but somehow her relatively stout body escaped your clutches. Within a few weeks, swarms of moths in your tattered wardrobe suggest that Mama Moth was in the family way.

There must be at least 200 critters nesting in every crevice of your clothing.

Two hundred plus moth-tykes from one female moth—is this possible? Indeed it is. The number of eggs, $f(x)$, in a female moth is a function of her abdominal width, x , in millimeters, modeled by

$$f(x) = 14x^3 - 17x^2 - 16x + 34, \quad 1.5 \leq x \leq 3.5.$$

Because there are 200 moths feasting on your favorite sweaters, Mama's abdominal width can be estimated by finding the solutions of the polynomial equation

$$14x^3 - 17x^2 - 16x + 34 = 200.$$

How can we solve such an equation? You might begin by subtracting 200 from both sides to obtain zero on one side. But then what? The factoring that we used in the previous section will not work in this situation.

In Section 2.5, we will present techniques for solving certain kinds of polynomial equations. These techniques will further enhance your ability to manipulate algebraically the polynomial functions that model your world. Because these techniques are based on understanding polynomial division, in this section we look at two methods for dividing polynomials. (We'll return to Mama Moth's abdominal width in the exercise set.)

- 1 Use long division to divide polynomials.

Long Division of Polynomials and the Division Algorithm

We begin by looking at division by a polynomial containing more than one term, such as

$$x + 3 \overline{)x^2 + 10x + 21}.$$

Divisor has two terms and is a binomial.

The polynomial dividend has three terms and is a trinomial.

When a divisor has more than one term, the four steps used to divide whole numbers—**divide, multiply, subtract, bring down the next term**—form the repetitive procedure for polynomial long division.

EXAMPLE 1 Long Division of Polynomials

Divide $x^2 + 10x + 21$ by $x + 3$.

Solution The following steps illustrate how polynomial division is very similar to numerical division.

$$x + 3 \overline{)x^2 + 10x + 21}$$

Arrange the terms of the dividend ($x^2 + 10x + 21$) and the divisor ($x + 3$) in descending powers of x .

$$x + 3 \overline{)x^2 + 10x + 21}$$

Divide x^2 (the first term in the dividend) by x (the first term in the divisor): $\frac{x^2}{x} = x$. Align like terms.

$$x(x + 3) = x^2 + 3x$$

$$x + 3 \overline{)x^2 + 10x + 21}$$

$$\underline{x^2 + 3x}$$

Multiply each term in the divisor ($x + 3$) by x , aligning terms of the product under like terms in the dividend.

$$\begin{array}{r}
 x \\
 x + 3 \overline{) x^2 + 10x + 21} \\
 \underline{x^2 + 3x} \\
 7x
 \end{array}$$

Change signs of the polynomial being subtracted.

Subtract $x^2 + 3x$ from $x^2 + 10x$ by changing the sign of each term in the lower expression and adding.

$$\begin{array}{r}
 x \\
 x + 3 \overline{) x^2 + 10x + 21} \\
 \underline{x^2 + 3x} \\
 7x + 21
 \end{array}$$

Bring down 21 from the original dividend and add algebraically to form a new dividend.

$$\begin{array}{r}
 x + 7 \\
 x + 3 \overline{) x^2 + 10x + 21} \\
 \underline{x^2 + 3x} \\
 7x + 21
 \end{array}$$

Find the second term of the quotient. Divide the first term of $7x + 21$ by x , the first term of the divisor: $\frac{7x}{x} = 7$.

$$\begin{array}{r}
 x + 7 \\
 x + 3 \overline{) x^2 + 10x + 21} \\
 \underline{x^2 + 3x} \\
 7x + 21 \\
 \underline{7x + 21} \\
 0
 \end{array}$$

$7(x + 3) = 7x + 21$

Remainder

Multiply the divisor $(x + 3)$ by 7, aligning under like terms in the new dividend. Then subtract to obtain the remainder of 0.

The quotient is $x + 7$. Because the remainder is 0, we can conclude that $x + 3$ is a factor of $x^2 + 10x + 21$ and

$$\frac{x^2 + 10x + 21}{x + 3} = x + 7.$$

 **Check Point** | Divide $x^2 + 14x + 45$ by $x + 9$.

Before considering additional examples, let's summarize the general procedure for dividing one polynomial by another.

Long Division of Polynomials

1. Arrange the terms of both the dividend and the divisor in descending powers of any variable.
2. **Divide** the first term in the dividend by the first term in the divisor. The result is the first term of the quotient.
3. **Multiply** every term in the divisor by the first term in the quotient. Write the resulting product beneath the dividend with like terms lined up.
4. **Subtract** the product from the dividend.
5. **Bring down** the next term in the original dividend and write it next to the remainder to form a new dividend.
6. Use this new expression as the dividend and repeat this process until the remainder can no longer be divided. This will occur when the degree of the remainder (the highest exponent on a variable in the remainder) is less than the degree of the divisor.

In our next long division, we will obtain a nonzero remainder.

EXAMPLE 2 Long Division of Polynomials

Divide $4 - 5x - x^2 + 6x^3$ by $3x - 2$.

Solution We begin by writing the dividend in descending powers of x .

$$4 - 5x - x^2 + 6x^3 = 6x^3 - x^2 - 5x + 4$$

Change signs of the polynomial being subtracted.

Divide: $\frac{6x^3}{3x} = 2x^2$.

Multiply: $2x^2(3x - 2) = 6x^3 - 4x^2$.

Subtract $6x^3 - 4x^2$ from $6x^3 - x^2$ and bring down $-5x$.

Now we divide $3x^2$ by $3x$ to obtain x , multiply x and the divisor, and subtract.

Change signs of the polynomial being subtracted.

Divide: $\frac{3x^2}{3x} = x$.

Multiply: $x(3x - 2) = 3x^2 - 2x$.

Subtract $3x^2 - 2x$ from $3x^2 - 5x$ and bring down 4 .

Now we divide $-3x$ by $3x$ to obtain -1 , multiply -1 and the divisor, and subtract.

Change signs of the polynomial being subtracted.

Divide: $\frac{-3x}{3x} = -1$.

Multiply: $-1(3x - 2) = -3x + 2$.

Subtract $-3x + 2$ from $-3x + 4$, leaving a remainder of 2 .

Remainder

The quotient is $2x^2 + x - 1$ and the remainder is 2 . When there is a nonzero remainder, as in this example, list the quotient, plus the remainder above the divisor. Thus,

$$\frac{6x^3 - x^2 - 5x + 4}{3x - 2} = \underbrace{2x^2 + x - 1}_{\text{Quotient}} + \frac{2}{3x - 2} \quad \text{Remainder above divisor}$$

An important property of division can be illustrated by clearing fractions in the equation that concluded Example 2. Multiplying both sides of this equation by $3x - 2$ results in the following equation:

$$6x^3 - x^2 - 5x + 4 = (3x - 2)(2x^2 + x - 1) + 2$$

Dividend
Divisor
Quotient
Remainder


Polynomial long division is checked by multiplying the divisor with the quotient and then adding the remainder. This should give the dividend. The process illustrates the **Division Algorithm**.

The Division Algorithm

If $f(x)$ and $d(x)$ are polynomials, with $d(x) \neq 0$, and the degree of $d(x)$ is less than or equal to the degree of $f(x)$, then there exist unique polynomials $q(x)$ and $r(x)$ such that

$$\begin{array}{ccccccc}
 f(x) & = & d(x) & \cdot & q(x) & + & r(x) \\
 \text{Dividend} & & \text{Divisor} & & \text{Quotient} & & \text{Remainder}
 \end{array}$$

The remainder, $r(x)$, equals 0 or it is of degree less than the degree of $d(x)$. If $r(x) = 0$, we say that $d(x)$ **divides evenly** into $f(x)$ and that $d(x)$ and $q(x)$ are **factors** of $f(x)$.

 **Check Point 2** Divide $7 - 11x - 3x^2 + 2x^3$ by $x - 3$. Express the result in the form quotient, plus remainder divided by divisor.

If a power of x is missing in either a dividend or a divisor, add that power of x with a coefficient of 0 and then divide. In this way, like terms will be aligned as you carry out the long division.

EXAMPLE 3 Long Division of Polynomials

Divide $6x^4 + 5x^3 + 3x - 5$ by $3x^2 - 2x$.

Solution We write the dividend, $6x^4 + 5x^3 + 3x - 5$, as $6x^4 + 5x^3 + 0x^2 + 3x - 5$ to keep all like terms aligned.


Multiply.

$$\begin{array}{r}
 2x^2 + 3x + 2 \\
 \hline
 3x^2 - 2x \overline{) 6x^4 + 5x^3 + 0x^2 + 3x - 5} \\
 \underline{\ominus 6x^4 - 4x^3} \\
 9x^3 + 0x^2 \\
 \underline{\ominus 9x^3 - 6x^2} \\
 6x^2 + 3x - 5 \\
 \underline{\ominus 6x^2 - 4x} \\
 7x - 5 \\
 \hline
 \text{Remainder}
 \end{array}$$

$2x^2(3x^2 - 2x) = 6x^4 - 4x^3$
 $3x(3x^2 - 2x) = 9x^3 - 6x^2$
 $2(3x^2 - 2x) = 6x^2 - 4x$

The division process is finished because the degree of $7x - 5$, which is 1, is less than the degree of the divisor $3x^2 - 2x$, which is 2. The answer is

$$\frac{6x^4 + 5x^3 + 3x - 5}{3x^2 - 2x} = 2x^2 + 3x + 2 + \frac{7x - 5}{3x^2 - 2x}$$

 **Check Point 3** Divide $2x^4 + 3x^3 - 7x - 10$ by $x^2 - 2x$.

2 Use synthetic division to divide polynomials.

Dividing Polynomials Using Synthetic Division

We can use **synthetic division** to divide polynomials if the divisor is of the form $x - c$. This method provides a quotient more quickly than long division. Let's compare the two methods showing $x^3 + 4x^2 - 5x + 5$ divided by $x - 3$.

Long Division

$$\begin{array}{r}
 \overline{) x^3 + 4x^2 - 5x + 5} \\
 \underline{\ominus x^3 - 3x^2} \\
 7x^2 - 5x + 5 \\
 \underline{\ominus 7x^2 - 21x} \\
 16x + 5 \\
 \underline{\ominus 16x - 48} \\
 53 \\
 \hline
 \text{Remainder}
 \end{array}$$

Divisor: $x - c$; $c = 3$
 Quotient: $x^2 + 7x + 16$
 Dividend: $x^3 + 4x^2 - 5x + 5$

Synthetic Division

$$\begin{array}{r|rrrr}
 3 & 1 & 4 & -5 & 5 \\
 & & 3 & 21 & 48 \\
 \hline
 & 1 & 7 & 16 & 53
 \end{array}$$

Long Division

$$\begin{array}{r}
 \overline{x^2 + 7x + 16} \\
 x-3 \overline{x^3 + 4x^2 - 5x + 5} \\
 \underline{x^3 - 3x^2} \\
 7x^2 - 5x \\
 \underline{7x^2 - 21x} \\
 16x + 5 \\
 \underline{16x - 48} \\
 53
 \end{array}$$

Quotient
Dividend
Divisor
 $x - c$;
 $c = 3$
Remainder

Notice the relationship between the polynomials in the long division process and the numbers that appear in synthetic division.

These are the coefficients of the dividend $x^3 + 4x^2 - 5x + 5$.

$$\begin{array}{r}
 3 \overline{1 \quad 4 \quad -5 \quad 5} \\
 \underline{3 \quad 21 \quad 48} \\
 1 \quad 7 \quad 16 \quad 53
 \end{array}$$

The divisor is $x - 3$. This is 3, or c , in $x - c$.

These are the coefficients of the quotient $x^2 + 7x + 16$.

This is the remainder.

Now let's look at the steps involved in synthetic division.

Synthetic Division

To divide a polynomial by $x - c$:

1. Arrange the polynomial in descending powers, with a 0 coefficient for any missing term.
2. Write c for the divisor, $x - c$. To the right, write the coefficients of the dividend.
3. Write the leading coefficient of the dividend on the bottom row.
4. Multiply c (in this case, 3) times the value just written on the bottom row. Write the product in the next column in the second row.
5. Add the values in this new column, writing the sum in the bottom row.
6. Repeat this series of multiplications and additions until all columns are filled in.
7. Use the numbers in the last row to write the quotient, plus the remainder above the divisor. **The degree of the first term of the quotient is one less than the degree of the first term of the dividend.** The final value in this row is the remainder.

Example

$$x - 3 \overline{x^3 + 4x^2 - 5x + 5}$$

$$3 \overline{1 \quad 4 \quad -5 \quad 5}$$

$$\begin{array}{r}
 3 \overline{1 \quad 4 \quad -5 \quad 5} \\
 \downarrow \text{Bring down 1.} \\
 1
 \end{array}$$

$$\begin{array}{r}
 3 \overline{1 \quad 4 \quad -5 \quad 5} \\
 \downarrow 3 \\
 1 \quad 3
 \end{array}$$

Multiply by 3: $3 \cdot 1 = 3$.

$$\begin{array}{r}
 3 \overline{1 \quad 4 \quad -5 \quad 5} \\
 \downarrow 3 \quad \text{Add.} \\
 1 \quad 7
 \end{array}$$

$$\begin{array}{r}
 3 \overline{1 \quad 4 \quad -5 \quad 5} \\
 \downarrow 3 \quad 21 \quad \text{Add.} \\
 1 \quad 7 \quad 16
 \end{array}$$

Multiply by 3: $3 \cdot 7 = 21$.

$$\begin{array}{r}
 3 \overline{1 \quad 4 \quad -5 \quad 5} \\
 \downarrow 3 \quad 21 \quad 48 \quad \text{Add.} \\
 1 \quad 7 \quad 16 \quad 53
 \end{array}$$

Multiply by 3: $3 \cdot 16 = 48$.

Written from the last row of the synthetic division

$$\begin{array}{r}
 1x^2 + 7x + 16 + \frac{53}{x - 3} \\
 x - 3 \overline{x^3 + 4x^2 - 5x + 5}
 \end{array}$$

EXAMPLE 4 Using Synthetic Division

Use synthetic division to divide $5x^3 + 6x + 8$ by $x + 2$.

Solution The divisor must be in the form $x - c$. Thus, we write $x + 2$ as $x - (-2)$. This means that $c = -2$. Writing a 0 coefficient for the missing x^2 -term in the dividend, we can express the division as follows:

$$x - (-2) \overline{)5x^3 + 0x^2 + 6x + 8.}$$

Now we are ready to set up the problem so that we can use synthetic division.

Use the coefficients of the dividend
 $5x^3 + 0x^2 + 6x + 8$ in descending powers of x .

This is c in
 $x - (-2)$.

$$\begin{array}{r|rrrr} -2 & 5 & 0 & 6 & 8 \end{array}$$

We begin the synthetic division process by bringing down 5. This is followed by a series of multiplications and additions.

1. Bring down 5.

$$\begin{array}{r|rrrr} -2 & 5 & 0 & 6 & 8 \\ & \downarrow & & & \\ & 5 & & & \end{array}$$

2. Multiply: $-2(5) = -10$.

$$\begin{array}{r|rrrr} -2 & 5 & 0 & 6 & 8 \\ & \downarrow & & & \\ & 5 & & & \\ & & -10 & & \end{array}$$

Multiply 5 by -2 .

3. Add: $0 + (-10) = -10$.

$$\begin{array}{r|rrrr} -2 & 5 & 0 & 6 & 8 \\ & \downarrow & & & \\ & 5 & -10 & & \\ & & & -10 & \end{array}$$

Add.

4. Multiply: $-2(-10) = 20$.

$$\begin{array}{r|rrrr} -2 & 5 & 0 & 6 & 8 \\ & \downarrow & & & \\ & 5 & -10 & & \\ & & & 20 & \\ & & 5 & -10 & \end{array}$$

Multiply -10 by -2 .

5. Add: $6 + 20 = 26$.

$$\begin{array}{r|rrrr} -2 & 5 & 0 & 6 & 8 \\ & \downarrow & & & \\ & 5 & -10 & 26 & \\ & & 5 & -10 & 26 \end{array}$$

Add.

6. Multiply: $-2(26) = -52$.

$$\begin{array}{r|rrrr} -2 & 5 & 0 & 6 & 8 \\ & \downarrow & & & \\ & 5 & -10 & 26 & \\ & & 5 & -10 & 26 \\ & & & -52 & \\ & & & & 26 \end{array}$$

Multiply 26 by -2 .

7. Add: $8 + (-52) = -44$.

$$\begin{array}{r|rrrr} -2 & 5 & 0 & 6 & 8 \\ & \downarrow & & & \\ & 5 & -10 & 26 & -52 \\ & & 5 & -10 & 26 \\ & & & -44 & \end{array}$$

Add.

The numbers in the last row represent the coefficients of the quotient and the remainder. The degree of the first term of the quotient is one less than that of the dividend. Because the degree of the dividend, $5x^3 + 6x + 8$, is 3, the degree of the quotient is 2. This means that the 5 in the last row represents $5x^2$.

$$\begin{array}{r|rrrr} -2 & 5 & 0 & 6 & 8 \\ & \downarrow & & & \\ & 5 & -10 & 26 & -52 \\ & & 5 & -10 & 26 \\ & & & -44 & \end{array}$$

The quotient is $5x^2 - 10x + 26$. The remainder is -44 .

Thus,

$$\begin{array}{r} 5x^2 - 10x + 26 - \frac{44}{x + 2} \\ x + 2 \overline{)5x^3 + 6x + 8} \end{array}$$

 **Check Point 4** Use synthetic division to divide $x^3 - 7x - 6$ by $x + 2$.

- 3 Evaluate a polynomial using the Remainder Theorem.

The Remainder Theorem

Let's consider the Division Algorithm when the dividend, $f(x)$, is divided by $x - c$. In this case, the remainder must be a constant because its degree is less than one, the degree of $x - c$.

$$f(x) = (x - c)q(x) + r$$

Dividend
Divisor
Quotient
Constant remainder

The remainder, r , is a constant when dividing by $x - c$.

Now let's evaluate f at c .

$$f(c) = (c - c)q(c) + r \quad \text{Find } f(c) \text{ by letting } x = c \text{ in } f(x) = (x - c)q(x) + r. \text{ This will give an expression for } r.$$

$$f(c) = 0 \cdot q(c) + r \quad c - c = 0$$

$$f(c) = r \quad 0 \cdot q(c) = 0 \text{ and } 0 + r = r.$$

What does this last equation mean? If a polynomial is divided by $x - c$, the remainder is the value of the polynomial at c . This result is called the **Remainder Theorem**.

The Remainder Theorem

If the polynomial $f(x)$ is divided by $x - c$, then the remainder is $f(c)$.

Example 5 shows how we can use the Remainder Theorem to evaluate a polynomial function at 2. Rather than substituting 2 for x , we divide the function by $x - 2$. The remainder is $f(2)$.

EXAMPLE 5 Using the Remainder Theorem to Evaluate a Polynomial Function

Given $f(x) = x^3 - 4x^2 + 5x + 3$, use the Remainder Theorem to find $f(2)$.

Solution By the Remainder Theorem, if $f(x)$ is divided by $x - 2$, then the remainder is $f(2)$. We'll use synthetic division to divide.

$$\begin{array}{r|rrrr} 2 & 1 & -4 & 5 & 3 \\ & & 2 & -4 & 2 \\ \hline & 1 & -2 & 1 & 5 \end{array}$$

Remainder

The remainder, 5, is the value of $f(2)$. Thus, $f(2) = 5$. We can verify that this is correct by evaluating $f(2)$ directly. Using $f(x) = x^3 - 4x^2 + 5x + 3$, we obtain

$$f(2) = 2^3 - 4 \cdot 2^2 + 5 \cdot 2 + 3 = 8 - 16 + 10 + 3 = 5. \quad \bullet$$

Check Point 5 Given $f(x) = 3x^3 + 4x^2 - 5x + 3$, use the Remainder Theorem to find $f(-4)$.

- 4 Use the Factor Theorem to solve a polynomial equation.

The Factor Theorem

Let's look again at the Division Algorithm when the divisor is of the form $x - c$.

$$f(x) = (x - c)q(x) + r$$

Dividend
Divisor
Quotient
Constant remainder

By the Remainder Theorem, the remainder r is $f(c)$, so we can substitute $f(c)$ for r :

$$f(x) = (x - c)q(x) + f(c).$$

Notice that if $f(c) = 0$, then

$$f(x) = (x - c)q(x)$$

so that $x - c$ is a factor of $f(x)$. This means that for the polynomial function $f(x)$, if $f(c) = 0$, then $x - c$ is a factor of $f(x)$.

Let's reverse directions and see what happens if $x - c$ is a factor of $f(x)$. This means that

$$f(x) = (x - c)q(x).$$

If we replace x in $f(x) = (x - c)q(x)$ with c , we obtain

$$f(c) = (c - c)q(c) = 0 \cdot q(c) = 0.$$

Thus, if $x - c$ is a factor of $f(x)$, then $f(c) = 0$.

We have proved a result known as the **Factor Theorem**.

The Factor Theorem

Let $f(x)$ be a polynomial.

- If $f(c) = 0$, then $x - c$ is a factor of $f(x)$.
- If $x - c$ is a factor of $f(x)$, then $f(c) = 0$.

The example that follows shows how the Factor Theorem can be used to solve a polynomial equation.

EXAMPLE 6 Using the Factor Theorem

Solve the equation $2x^3 - 3x^2 - 11x + 6 = 0$ given that 3 is a zero of $f(x) = 2x^3 - 3x^2 - 11x + 6$.

Solution We are given that 3 is a zero of $f(x) = 2x^3 - 3x^2 - 11x + 6$. This means that $f(3) = 0$. Because $f(3) = 0$, the Factor Theorem tells us that $x - 3$ is a factor of $f(x)$. We'll use synthetic division to divide $f(x)$ by $x - 3$.

$$\begin{array}{r|rrrr} 3 & 2 & -3 & -11 & 6 \\ & & 6 & 9 & -6 \\ \hline & 2 & 3 & -2 & 0 \end{array} \quad x - 3 \overline{) 2x^3 - 3x^2 - 11x + 6}$$

Equivalently,

$$2x^3 - 3x^2 - 11x + 6 = (x - 3)(2x^2 + 3x - 2).$$

The remainder, 0, verifies that $x - 3$ is a factor of $2x^3 - 3x^2 - 11x + 6$.

Now we can solve the polynomial equation.

$$\begin{aligned} 2x^3 - 3x^2 - 11x + 6 &= 0 && \text{This is the given equation.} \\ (x - 3)(2x^2 + 3x - 2) &= 0 && \text{Factor using the result from the synthetic division.} \\ (x - 3)(2x - 1)(x + 2) &= 0 && \text{Factor the trinomial.} \\ x - 3 = 0 \quad \text{or} \quad 2x - 1 = 0 \quad \text{or} \quad x + 2 = 0 &&& \text{Set each factor equal to 0.} \\ x = 3 \quad \quad \quad x = \frac{1}{2} \quad \quad \quad x = -2 &&& \text{Solve for } x. \end{aligned}$$

The solution set is $\{-2, \frac{1}{2}, 3\}$.

Based on the Factor Theorem, the following statements are useful in solving polynomial equations:

- If $f(x)$ is divided by $x - c$ and the remainder is zero, then c is a zero of f and c is a root of the polynomial equation $f(x) = 0$.
- If $f(x)$ is divided by $x - c$ and the remainder is zero, then $x - c$ is a factor of $f(x)$.

Technology

Graphic Connections


Because the solution set of

$$2x^3 - 3x^2 - 11x + 6 = 0$$

is $\{-2, \frac{1}{2}, 3\}$, this implies that the polynomial function

$$f(x) = 2x^3 - 3x^2 - 11x + 6$$

has x -intercepts (or zeros) at $-2, \frac{1}{2}$, and 3. This is verified by the graph of f .



Check Point 6 Solve the equation $15x^3 + 14x^2 - 3x - 2 = 0$ given that -1 is a zero of $f(x) = 15x^3 + 14x^2 - 3x - 2$.

Exercise Set 2.4

Practice Exercises

In Exercises 1–16, divide using long division. State the quotient, $q(x)$, and the remainder, $r(x)$.

1. $(x^2 + 8x + 15) \div (x + 5)$
2. $(x^2 + 3x - 10) \div (x - 2)$
3. $(x^3 + 5x^2 + 7x + 2) \div (x + 2)$
4. $(x^3 - 2x^2 - 5x + 6) \div (x - 3)$
5. $(6x^3 + 7x^2 + 12x - 5) \div (3x - 1)$
6. $(6x^3 + 17x^2 + 27x + 20) \div (3x + 4)$
7. $(12x^2 + x - 4) \div (3x - 2)$
8. $(4x^2 - 8x + 6) \div (2x - 1)$
9. $\frac{2x^3 + 7x^2 + 9x - 20}{x + 3}$
10. $\frac{3x^2 - 2x + 5}{x - 3}$
11. $\frac{4x^4 - 4x^2 + 6x}{x - 4}$
12. $\frac{x^4 - 81}{x - 3}$
13. $\frac{6x^3 + 13x^2 - 11x - 15}{3x^2 - x - 3}$
14. $\frac{x^4 + 2x^3 - 4x^2 - 5x - 6}{x^2 + x - 2}$
15. $\frac{18x^4 + 9x^3 + 3x^2}{3x^2 + 1}$
16. $\frac{2x^5 - 8x^4 + 2x^3 + x^2}{2x^3 + 1}$

In Exercises 17–32, divide using synthetic division.

17. $(2x^2 + x - 10) \div (x - 2)$
18. $(x^2 + x - 2) \div (x - 1)$
19. $(3x^2 + 7x - 20) \div (x + 5)$
20. $(5x^2 - 12x - 8) \div (x + 3)$
21. $(4x^3 - 3x^2 + 3x - 1) \div (x - 1)$
22. $(5x^3 - 6x^2 + 3x + 11) \div (x - 2)$
23. $(6x^5 - 2x^3 + 4x^2 - 3x + 1) \div (x - 2)$
24. $(x^5 + 4x^4 - 3x^2 + 2x + 3) \div (x - 3)$
25. $(x^2 - 5x - 5x^3 + x^4) \div (5 + x)$
26. $(x^2 - 6x - 6x^3 + x^4) \div (6 + x)$
27. $\frac{x^5 + x^3 - 2}{x - 1}$
28. $\frac{x^7 + x^5 - 10x^3 + 12}{x + 2}$
29. $\frac{x^4 - 256}{x - 4}$
30. $\frac{x^7 - 128}{x - 2}$
31. $\frac{2x^5 - 3x^4 + x^3 - x^2 + 2x - 1}{x + 2}$
32. $\frac{x^5 - 2x^4 - x^3 + 3x^2 - x + 1}{x - 2}$

In Exercises 33–40, use synthetic division and the Remainder Theorem to find the indicated function value.

33. $f(x) = 2x^3 - 11x^2 + 7x - 5$; $f(4)$
34. $f(x) = x^3 - 7x^2 + 5x - 6$; $f(3)$
35. $f(x) = 3x^3 - 7x^2 - 2x + 5$; $f(-3)$
36. $f(x) = 4x^3 + 5x^2 - 6x - 4$; $f(-2)$
37. $f(x) = x^4 + 5x^3 + 5x^2 - 5x - 6$; $f(3)$
38. $f(x) = x^4 - 5x^3 + 5x^2 + 5x - 6$; $f(2)$
39. $f(x) = 2x^4 - 5x^3 - x^2 + 3x + 2$; $f\left(-\frac{1}{2}\right)$

40. $f(x) = 6x^4 + 10x^3 + 5x^2 + x + 1$; $f\left(-\frac{2}{3}\right)$

41. Use synthetic division to divide

$$f(x) = x^3 - 4x^2 + x + 6 \text{ by } x + 1.$$

Use the result to find all zeros of f .

42. Use synthetic division to divide

$$f(x) = x^3 - 2x^2 - x + 2 \text{ by } x + 1.$$


Use the result to find all zeros of f .

43. Solve the equation $2x^3 - 5x^2 + x + 2 = 0$ given that 2 is a zero of $f(x) = 2x^3 - 5x^2 + x + 2$.
44. Solve the equation $2x^3 - 3x^2 - 11x + 6 = 0$ given that -2 is a zero of $f(x) = 2x^3 - 3x^2 - 11x + 6$.
45. Solve the equation $12x^3 + 16x^2 - 5x - 3 = 0$ given that $-\frac{3}{2}$ is a root.
46. Solve the equation $3x^3 + 7x^2 - 22x - 8 = 0$ given that $-\frac{1}{3}$ is a root.


Practice Plus

In Exercises 47–50, use the graph or the table to determine a solution of each equation. Use synthetic division to verify that this number is a solution of the equation. Then solve the polynomial equation.

47. $x^3 + 2x^2 - 5x - 6 = 0$



48. $2x^3 + x^2 - 13x + 6 = 0$



49. $6x^3 - 11x^2 + 6x - 1 = 0$

$y_1 = 6x^3 - 11x^2 + 6x - 1$

X	Y1
-3	-280
-2	-105
-1	-24
0	-1
1	0
2	15
3	80

$X = -3$

50. $2x^3 + 11x^2 - 7x - 6 = 0$

$y_1 = 2x^3 + 11x^2 - 7x - 6$

X	Y ₁
3	60
-2	36
-1	10
0	-6
1	0
2	40
3	126

$X = -3$

Application Exercises

51. a. Use synthetic division to show that 3 is a solution of the polynomial equation

$$14x^3 - 17x^2 - 16x - 177 = 0.$$

- b. Use the solution from part (a) to solve this problem. The number of eggs, $f(x)$, in a female moth is a function of her abdominal width, x , in millimeters, modeled by


$$f(x) = 14x^3 - 17x^2 - 16x + 34.$$

What is the abdominal width when there are 211 eggs?

52. a. Use synthetic division to show that 2 is a solution of the polynomial equation

$$2h^3 + 14h^2 - 72 = 0.$$

- b. Use the solution from part (a) to solve this problem. The width of a rectangular box is twice the height and the length is 7 inches more than the height. If the volume is 72 cubic inches, find the dimensions of the box.



In Exercises 53–54, write a polynomial that represents the length of each rectangle.


53. The width is $x + 0.2$ units. The area is $0.5x^3 - 0.3x^2 + 0.22x + 0.06$ square units.

54. The width is $x + \frac{3}{4}$ units. The area is $8x^3 - 6x^2 - 5x + 3$ square units.

During the 1980s, the controversial economist Arthur Laffer promoted the idea that tax increases lead to a reduction in government revenue. Called *supply-side economics*, the theory uses functions such as

$$f(x) = \frac{80x - 8000}{x - 110}, 30 \leq x \leq 100.$$

This function models the government tax revenue, $f(x)$, in tens of billions of dollars, in terms of the tax rate, x . The graph of the function is shown. It illustrates tax revenue decreasing quite dramatically as the tax rate increases. At a tax rate of (gasp) 100%, the government takes all our money and no one has an incentive to work. With no income earned, zero dollars in tax revenue is generated.



Use function f and its graph to solve Exercises 55–56.

55. a. Find and interpret $f(30)$. Identify the solution as a point on the graph of the function.

- b. Rewrite the function by using long division to perform $(80x - 8000) \div (x - 110)$.

Then use this new form of the function to find $f(30)$. Do you obtain the same answer as you did in part (a)?

- c. Is f a polynomial function? Explain your answer.

56. a. Find and interpret $f(40)$. Identify the solution as a point on the graph of the function.

- b. Rewrite the function by using long division to perform $(80x - 8000) \div (x - 110)$.

Then use this new form of the function to find $f(40)$. Do you obtain the same answer as you did in part (a)?

- c. Is f a polynomial function? Explain your answer.

Writing in Mathematics

57. Explain how to perform long division of polynomials. Use $2x^3 - 3x^2 - 11x + 7$ divided by $x - 3$ in your explanation.

58. In your own words, state the Division Algorithm.

59. How can the Division Algorithm be used to check the quotient and remainder in a long division problem?

60. Explain how to perform synthetic division. Use the division problem in Exercise 57 to support your explanation.

61. State the Remainder Theorem.

62. Explain how the Remainder Theorem can be used to find $f(-6)$ if $f(x) = x^4 + 7x^3 + 8x^2 + 11x + 5$. What advantage is there to using the Remainder Theorem in this situation rather than evaluating $f(-6)$ directly?

63. How can the Factor Theorem be used to determine if $x - 1$ is a factor of $x^3 - 2x^2 - 11x + 12$?

64. If you know that -2 is a zero of

$$f(x) = x^3 + 7x^2 + 4x - 12,$$

explain how to solve the equation

$$x^3 + 7x^2 + 4x - 12 = 0.$$

Technology Exercise

65. For each equation that you solved in Exercises 43–46, use a graphing utility to graph the polynomial function defined by the left side of the equation. Use end behavior to obtain a complete graph. Then use the graph's x -intercepts to verify your solutions.

Critical Thinking Exercises

Make Sense? In Exercises 66–69, determine whether each statement makes sense or does not make sense, and explain your reasoning.

66. When performing the division $(x^5 + 1) \div (x + 1)$, there's no need for me to follow all the steps involved in polynomial long division because I can work the problem in my head and see that the quotient must be $x^4 + 1$.
67. Every time I divide polynomials using synthetic division, I am using a highly condensed form of the long division procedure where omitting the variables and exponents does not involve the loss of any essential data.
68. The only nongraphic method that I have for evaluating a function at a given value is to substitute that value into the function's equation.
69. I found the zeros of function f , but I still need to find the solutions of the equation $f(x) = 0$.

In Exercises 70–73, determine whether each statement is true or false. If the statement is false, make the necessary change(s) to produce a true statement.

70. If a trinomial in x of degree 6 is divided by a trinomial in x of degree 3, the degree of the quotient is 2.

71. Synthetic division can be used to find the quotient of $10x^3 - 6x^2 + 4x - 1$ and $x - \frac{1}{2}$.
72. Any problem that can be done by synthetic division can also be done by the method for long division of polynomials.
73. If a polynomial long-division problem results in a remainder that is a whole number, then the divisor is a factor of the dividend.
74. Find k so that $4x + 3$ is a factor of

$$20x^3 + 23x^2 - 10x + k.$$

75. When $2x^2 - 7x + 9$ is divided by a polynomial, the quotient is $2x - 3$ and the remainder is 3. Find the polynomial.
76. Find the quotient of $x^{3n} + 1$ and $x^n + 1$.
77. Synthetic division is a process for dividing a polynomial by $x - c$. The coefficient of x in the divisor is 1. How might synthetic division be used if you are dividing by $2x - 4$?
78. Use synthetic division to show that 5 is a solution of

$$x^4 - 4x^3 - 9x^2 + 16x + 20 = 0.$$

Then solve the polynomial equation.

Preview Exercises

Exercises 79–81 will help you prepare for the material covered in the next section.

79. Solve: $x^2 + 4x - 1 = 0$.
80. Solve: $x^2 + 4x + 6 = 0$.
81. Let $f(x) = a_n(x^4 - 3x^2 - 4)$. If $f(3) = -150$, determine the value of a_n .

Section 2.5 Zeros of Polynomial Functions

Objectives

- 1 Use the Rational Zero Theorem to find possible rational zeros.
- 2 Find zeros of a polynomial function.
- 3 Solve polynomial equations.
- 4 Use the Linear Factorization Theorem to find polynomials with given zeros.
- 5 Use Descartes's Rule of Signs.



You stole my formula!

Tartaglia (1499–1557), poor and starving, has found a formula that gives a root for a third-degree polynomial equation. Cardano (1501–1576) begs Tartaglia to reveal the secret formula, wheedling it from him with the promise he will find the impoverished Tartaglia a patron. Then Cardano publishes his famous work *Ars Magna*, in which he presents Tartaglia's formula as his own. Cardano uses his most talented student, Ferrari (1522–1565), who derived a formula for a root of a fourth-

Tartaglia's Secret Formula for One Solution of $x^3 + mx = n$

$$x = \sqrt[3]{\sqrt{\left(\frac{n}{2}\right)^2 + \left(\frac{m}{3}\right)^3} + \frac{n}{2}} - \sqrt[3]{\sqrt{\left(\frac{n}{2}\right)^2 + \left(\frac{m}{3}\right)^3} - \frac{n}{2}}$$

Popularizers of mathematics are sharing bizarre stories that are giving math a secure place in popular culture. One episode, able to compete with the wildest fare served up by television talk shows and the tabloids, involves three Italian mathematicians and, of all things, zeros of polynomial functions.